## NTP Technical Report on Toxicity Studies of

## **Black Newsprint Inks**

Administered Topically to F344/N Rats and C3H Mice

Joel F. Mahler, DVM, Study Scientist National Toxicology Program Post Office Box 12233 Research Triangle Park, NC 27709

> NIH Publication 92-3340 July 1992

United States Department of Health and Human Services
Public Health Service
National Institutes of Health

### **CONTRIBUTORS**

The NTP report on the toxicity studies of black newsprint ink is based primarily on 30-day and 13-week studies performed over the period of December, 1987, to September, 1988, at Battelle Memorial Laboratories, Columbus Division, Columbus, Ohio.

#### **National Toxicology Program**

Evaluated experiment, interpreted results, and reported findings

Joel F. Mahler, DVM Study Scientist John R. Bucher, PhD Leo T. Burka, PhD Rajendra S. Chhabra, PhD Michael P. Dieter, PhD Michael R. Elwell, DVM, PhD H.B. Matthews, PhD Morrow Thompson, DVM, PhD Errol Zeiger, PhD

Coordinated report preparation

Jane M. Lambert, BS Kristine Witt, MS Oak Ridge Associated Universities

#### Battelle Memorial Laboratories, Columbus, OH

Principal contributors

Arthur C. Peters, DVM
Principal Investigator
Milton Hejtmancik, PhD
Study Director, Toxicologist
Ming Chang, PhD
Chemistry
Peter Jepsen, DVM, MS
Study Veterinarian
Ronald Pershing, DVM
Study Pathologist
Michael Ryan, DVM
Clinical Pathology

#### NTP Pathology Working Group

Evaluated slides and prepared pathology report

John Seely, DVM
Chairperson
Pathco, Inc.
Michael R. Elwell, DVM, PhD
National Toxicology Program

## Experimental Pathology Laboratories, Inc.

Provided pathology quality assessment

Cynthia Shackelford, PhD William MacKenzie, DVM, MS

# Environmental Health Research and Testing Inc.

Sperm morphology and vaginal cytology evaluation

Dushant K. Gulati, PhD Teresa Cocanougher, BA Susan Russell, BA

#### Analytical Sciences, Inc.

Statistical analysis

Steven Seilkop, MS Janet Teague, MS

# TABLE OF CONTENTS

CONTRIB	UTORS	2
TABLE OF	CONTENTS	3
ABSTRAC	т	5
PEER REV	VIEW PANEL	7
SUMMAR	Y OF PEER REVIEW COMMENTS	8
INTRODU	CTION	9
Dhygi	cal Properties, Production, and Exposure	C
Enide	miology	
	al Studies	
	genicity	
•	Rationale and Design	
MATERIA	LS AND METHODS	12
Procu	rement and Characterization of Samples	12
	y Study Design	
13-We	eek Study Design	13
	ductive Toxicity	
	ic Toxicology	
	tical Methods	
Quali	ty Assurance	16
RESULTS		18
	y Studies in F344/N Rats	
13-We	eek Studies in F344/N Rats	18
30-Da	y Studies in C3H Mice	21
13-We	eek Studies in C3H Mice	22
Genet	ic Toxicology	31
Discussi	ON	32
REFEREN	CES	34
TABLES		
	Emperimental Design and Materials and Mathada in the 20 Des	
Table 1	Experimental Design and Materials and Methods in the 30-Day and 13-Week Dermal Studies of Black Newsprint Inks	17
Table 2	Survival and Weight Gain of F344/N Rats in the 30-Day Dermal Studies	
	of Black Newsprint Inks	19
Table 3	Survival and Weight Gain of F344/N Rats in the 13-Week Dermal Studies	10
	of Black Newsprint Inks	19
Table 4	Selected Organ Weights of F344/N Rats in the 13-Week Dermal Studies	
	of Black Newsprint Inks	21
Table 5	Survival and Weight Gain of C3H Mice in the 30-Day Dermal Studies	
	of Black Newsprint Inks	22

Table 6 Survival and Weight Gain of C3H Mice in the 13-Week Dermal Studies of Black Newsprint Inks	4	BLACK NEWSPRINT INK, NTP TOXICITY REPORT NUMBER 17
of Black Newsprint Inks	Table 6	•
in the 13-Week Dermal Studies of Black Newsprint Inks in C3H Mice	Table 7	
Figure 1 Growth Curves of F344/N Rats in the 13-Week Dermal Studies of Black Newsprint Inks and Mineral Oils	Table 8	
of Black Newsprint Inks and Mineral Oils	FIGURES	
of Black Newsprint Inks and Mineral Oils	Figure 1	
APPENDICES  Appendix A Organ Weights and Organ-Weight-to-Body-Weight Ratios	Figure 2	
Appendix A Organ Weights and Organ-Weight-to-Body-Weight Ratios	PLATES	
Appendix B Hematology Results	APPENDICE	s
Appendix C Reproductive Tissue Evaluations and Estrous Cycle Length	Appendix A	Organ Weights and Organ-Weight-to-Body-Weight Ratios A-1
Appendix D Genetic ToxicologyD-1	Appendix B	Hematology ResultsB-1
	Appendix C	Reproductive Tissue Evaluations and Estrous Cycle Length
Appendix E Analyses for Polycyclic Aromatic Hydrocarbons E-1	Appendix D	Genetic ToxicologyD-1
	Appendix E	Analyses for Polycyclic Aromatic Hydrocarbons E-1

### **A**BSTRACT

Toxicity studies were conducted by applying black newsprint inks or mineral oils to clipped skin of the dorsal interscapular area of C3H mice and F344/N rats of both sexes, to determine systemic and local effects. Four lots of both letterpress and offset types of newsprint ink were studied, either as composite mixtures or as individual lots. An industrial grade mineral oil, used as an extender for newsprint ink formulation, and USP medicinal grade mineral oil also were studied. Analyses for the presence of polycyclic aromatic hydrocarbons (PAHs) were conducted on composite ink mixtures and mineral oils; letterpress and offset ink mixtures were found to have cumulative concentrations of 206 and 105 ppm, respectively; the concentration of PAHs in the printing ink mineral oil sample was 208 ppm, while none were detected in the USP grade mineral oil. In genetic toxicity studies, letterpress and offset newsprint ink composite mixtures were each mutagenic in *Salmonella typhimurium* strains TA98 and TA100 when tested in a preincubation protocol with added hamster liver S9. With rat liver S9, results for both inks were positive in strain TA98 and negative in strain TA100. Neither type of ink was mutagenic in the absence of S9 activation.

In 30-day studies, 5 rats and mice per sex were given single, daily dermal applications of letterpress or offset newsprint inks, 5 days per week, for a total of 21 - 22 applications. Dose groups for each type of ink received either the neat (undiluted) composite ink mixture, or the 3:1, 1:1, or 1:3 dilutions (ink:USP mineral oil), with a total dose volume of 100 (mice) or 250 (rats) µl. All animals survived until the end of the studies. Toxicity attributed to ink administration was limited to decreased body weight gains in female rats treated with neat and the 3:1 dilution of letterpress ink, and to scaliness at the site of application in 1 or more mice in each letterpress ink treatment group. As a result of grooming activity and the large amount of test chemical applied, chemicals were spread over the body, and there was evidence that some oral ingestion had occurred.

In 13-week studies, various ink and mineral oil formulations were administered dermally to 10 rats and mice per sex. To prevent accumulation of inks and distribution over the body as seen in the 30-day studies, the frequency of application was reduced to twice weekly and the total dose volume was decreased to 20 microliters for mice and 50 microliters for rats. Treatment groups for rats consisted of letterpress ink mixture, offset ink mixture, printing ink mineral oil, USP mineral oil, and clipped, untreated controls. Groups of mice were administered each of the 4 individual lots of both letterpress and offset inks, the composite mixtures of each, and printing ink and USP mineral oils; clipped, untreated groups served as controls. All rats, all male mice, and all female mice except one administered offset ink-lot E survived to the end of the studies. Effects attributable to compound administration in rats were limited to decreased body weight gains in females treated with printing ink mineral oil and letterpress ink mixture, and increased liver and kidney weights in both males and females exposed to USP mineral oil; there were no local toxic effects at the site of application. In mice, there were no body weight effects, but liver weights were increased in most ink and mineral oil treatment groups of both Dermal toxicity was evidenced in mice by scaliness and irritation at the site of sexes. application of both sexes treated with USP mineral oil and letterpress ink-lot C.

Microscopically, local toxicity at the site of application was observed in mice of all treatment groups and was characterized by acanthosis and inflammation.

In summary, results of these studies indicate that topical administration of black newsprint inks and mineral oils produces local toxicity at the site of application in mice; toxic effects on the skin in this species are consistent with those of a primary cutaneous irritant. In rats, possible evidence for toxicity was limited to decreased body weight gains in females treated with letterpress ink formulations.

### PEER REVIEW

#### **Peer Review Panel**

The members of the peer review panel who evaluated the draft report on the toxicity studies on black newsprint inks on July 10, 1991, are listed below. Panel members serve as independent scientists, not as representatives of any institution, company, or governmental agency. In this capacity, panel members act to determine if the design and conditions of the NTP studies were appropriate and to ensure that the toxicity study report fully and clearly presents the experimental results and conclusions.

### National Toxicology Program's Board of Scientific Counselors Technical Reports Review Subcommittee

Paul T. Bailey, PhD Mobil Oil Corporation Toxicology Division Princeton, NJ

Louis S. Beliczky, MS, MPH Director of Industrial Hygiene Department of Industrial Hygiene United Rubber Workers Intl. Union 87 South High Street Akron, OH

Gary P. Carlson, PhD Department of Pharmacology and Toxicology Purdue University West Lafayette, IN

Harold Davis, DVM, PhD School of Aerospace Medicine Brooks Air Force Base, TX

Robert H. Garman, DVM Consultants in Veterinary Pathology Murrysville, PA

Jay I. Goodman, PhD Department of Pharmacology and Toxicology Michigan State University East Lansing, MI David W. Hayden, DVM, PhD Department of Veterinary Pathobiology College of Veterinary Medicine University of Minnesota St. Paul, MN

Curtis D. Klaassen, PhD Department of Pharmacology and Toxicology University of Kansas Medical Center Kansas City, KS

Daniel S. Longnecker, MD Department of Pathology Dartmouth Medical School Hanover, NH

Barbara McKnight, PhD Department of Biostatistics University of Washington Seattle, WA

Ellen K. Silbergeld, PhD\* University of Maryland Medical School Baltimore, MD

Lauren Zeise, PhD California Department of Health Services Berkeley, CA

<sup>\*</sup>unable to attend

#### **Summary of Peer Review Comments**

On July 9 and 10, 1991, the Technical Reports Review Subcommittee of the Board of Scientific Counselors for the National Toxicology Program met in Research Triangle Park, NC, to review the draft technical report on toxicity studies of black newsprint inks.

Dr. J.F. Mahler, NIEHS, introduced the short-term toxicity studies of black newsprint inks by reviewing the uses of and rationale for study the inks, their composition, the methodology for topical application of the inks and mineral oils, experimental design, and results.

Dr. Davis, a principal reviewer, said the study was well-designed, very complex, and had been concisely and clearly reported. He said that the report summary mentioned little if any toxicity beyond the site of topical application, yet female rats had decreased weight gains and decreased heart and lung weights, compared to controls. He asked if these effects might be attributable to decreased feed consumption. Dr. Mahler agreed that the body-weight decreases were substantial enough to be seen as evidence of toxicity and that this would be noted in the report.

Dr. Bailey, a second principal reviewer, said the experiments were well-done and the report well-written. He cited references to the work of Blackburn *et al.* on the modified Ames assay and said these references should be included in the report, since the assay can accurately differentiate between carcinogenic and non-carcinogenic mineral oils and would also provide highly reliable estimates of the relative potency of such materials in mouse skin painting bioassays. Dr. Mahler said a later reference to the modified Ames assay was cited in the report, but that the Blackburn *et al.* work would be included as well.

Dr. Bailey said it would be helpful to include the refining histories for the printing ink mineral oils used in the study, since reference was made to the fact that less-refined oils have greater carcinogenic potential. Dr. Mahler said he would seek to include the refining histories of these oils in an appendix; however, subsequent efforts to locate further information concerning the refining histories were unsuccessful.

Dr. Bailey asked why the C3H mouse was used in the study. Dr. J. Bucher, NIEHS, noted that the National Institute of Occupational Safety and Health (NIOSH) had suggested use of that strain, saying it had a data base on dermal application studies using that type of animal. However, Dr. J. Haartz, NIOSH, said this database was not large and that much of it was based on studies involving asphalt fumes and fume fractions.

Mr. Beliczky said reference should be made to the fact that carbon black, used in some black newsprint inks, contains polycyclic aromatic hydrocarbons (PAHs); he noted that the International Agency for Research on Cancer (IARC) lists some PAHs in oil as human skin carcinogens. Dr. Mahler said that references cited in the report confirm Mr. Beliczky's comments.

After discussion of editorial matters, the panel agreed to accept the report, with the indicated changes.